

77538
Unusual Fragmental Breccia
47.2 grams



*Figure 1: Photo of 77538. Edge of 1 cm cube at bottom.
NASA S73-19065.*

Introduction

77538 is a trace-element-rich breccia sample. It is a fragmental breccia with a light-colored matrix (figure 2) It has patina and zap pits on all sides.

Petrography

Warner et al. (1977, 1978) reported that the olivine and some of the pyroxene in 77538 were Fe-rich (figure 4).

Taylor et al. (1980) reported granite clasts and suggested an origin by silicate-liquid immisibility.

Mineralogy

Olivine: Olivine is minor, but Fe-rich

Pyroxene: Some pyroxene is very iron rich (figure 4).

Plagioclase: Most plagioclase is An₈₅₋₉₈, but some is An₆₂.

Significant Clasts:

Granite clasts with intergrown K-spar and silica are present (figure 3).

Chemistry

Laul and Schmitt (1975) reported the composition in an abstract to the LPSC (table 1, figure 5). They found this sample had high Fe/Mg ratio and was very Zr - rich.

Processing

There are 6 thin sections.




Figure 2: Photomicrographs of thin section 77538, 17 by C.Meyer.




Figure 3: Thin section photomicrograph of "granite" clast in 77538 (from Warner et al. 1978). Field of view about 2 mm.




Figure 4: Pyroxene, olivine and plagioclase diagram for 77538 (from Warner et al. 1978).




Figure 5: Normalized rare-earth-element diagram for 77538 (data from Laul and Schmitt 1975).

Table 1. Chemical composition of 77538.

reference	Warner 77	Laul 75	Warner 78	Dalrymple96
weight				
SiO ₂ %		50.3	74	(b) 47.2 (c)
TiO ₂	1.2	1.2	(a) 2.82	(b) 1.6 (c)
Al ₂ O ₃	14.5	14.5	(a) 0.96	(b) 17.4 (c)
FeO	10.6	10.6	(a) 31.3	(b) 10.6 (a)
MnO	0.15	0.15	(a)	0.13 (c)
MgO	5	5	(a) 4.3	(b) 10.7 (c)
CaO	10.3	10.3	(a) 9.8	(b) 10.7 (c)
Na ₂ O	0.75	0.75	(a) 0.12	(b) 0.68 (a)
K ₂ O	1.04	1.04	(a) 0.29	(b) 0.25 (a)
P ₂ O ₅			0.28	0.07 (b)
S %				
sum				
Sc ppm		22	(a)	18.9 (a)
V		40	(a)	
Cr		1642	(a) 1163	(b) 1642 (a)
Co		13.5	(a)	38 (a)
Ni				314 (a)
Cu				
Zn				
Ga				
Ge ppb				
As				
Se				
Rb				10 (a)
Sr				176 (a)
Y				
Zr		730	(a)	530 (a)
Nb				
Mo				
Ru				
Rh				
Pd ppb				
Ag ppb				
Cd ppb				
In ppb				
Sn ppb				
Sb ppb				
Te ppb				
Cs ppm				0.36 (a)
Ba		700	(a)	434 (a)
La		59	(a)	40.4 (a)
Ce		150	(a)	104.2 (a)
Pr				
Nd		90	(a)	69 (a)
Sm		25.1	(a)	18.3 (a)
Eu		1.7	(a)	1.89 (a)
Gd				
Tb		5.5	(a)	3.8 (a)
Dy		35	(a)	
Ho				
Er				
Tm				
Yb		21.7	(a)	13.4 (a)
Lu		3.2	(a)	1.9 (a)
Hf		21.5	(a)	14.5 (a)
Ta		3.3	(a)	1.77 (a)
W ppb				
Re ppb				
Os ppb				
Ir ppb				8.8 (a)
Pt ppb				9.8 (a)
Au ppb				
Th ppm		16	(a)	7.1 (a)
U ppm		4.2	(a)	1.88 (a)

technique: (a) INAA, (b) broad beam electron probe,

References for 77538

Butler P. (1973) **Lunar Sample Information Catalog Apollo 17.** Lunar Receiving Laboratory. MSC 03211 Curator's Catalog. pp. 447.

Laul J.C. and Schmitt R.A. (1975c) Chemical composition of Apollo 17 samples: Boulder breccias (2), rake breccias (8), and others (abs). *Lunar Sci. VI*, 489-491. Lunar Planetary Institute, Houston

Meyer C. (1994) Catalog of Apollo 17 rocks. Vol. 4 North Massif

Muehlberger et al. (1973) Documentation and environment of the Apollo 17 samples: A preliminary report. Astrogeology 71 322 pp superceeded by Astrogeology 73 (1975) and by Wolfe et al. (1981)

Muehlberger W.R. and many others (1973) Preliminary Geological Investigation of the Apollo 17 Landing Site. In **Apollo 17 Preliminary Science Report.** NASA SP-330.

Taylor G.J., Warner R.D., Keil K., Ma M.-S. and Schmitt R.A. (1980) Silicate liquid immiscibility, evolved lunar rocks, and the formation of KREEP. In Proc. Conf. Lunar Highlands Crust, Geochim. Cosmochim. Acta Suppl. 12. Pergamon Press. 339-352. Lunar Planetary Institute, Houston.

Warner R.D., Keil K., Nehru C.E. and Taylor G.J. (1978a) Catalogue of Apollo 17 rake samples from Stations 1a, 2, 7, and 8. Spec. Publ. #18, UNM Institute of Meteoritics, Albuquerque. 88 pp.

Warner R.D., Taylor G.J., Mansker W.L. and Keil K. (1978b) Clast assemblages of possible deep-seated (77517) and immiscible melt (77538) origins in Apollo 17 breccias. Proc. 9th Lunar Planet. Sci. Conf. 941-958.

Warner R.D., Taylor G.J. and Keil K. (1978d) Clasts in breccias 77517 and 77538: Evidence for deep-seated and immiscible melt origins (abs). *Lunar Planet. Sci. IX*, 1222-1224. Lunar Planetary Institute, Houston.

Wolfe E.W., Bailey N.G., Lucchitta B.K., Muehlberger W.R., Scott D.H., Sutton R.L and Wilshire H.G. (1981) The geologic investigation of the Taurus-Littrow Valley: Apollo 17 Landing Site. US Geol. Survey Prof. Paper, 1080, pp. 280.

